

Abstract

Normal abundant dietary sugars such as fructose and glucose can contribute to hypertension and other health issues. To avoid these health complications, many individuals use artificial sweeteners. An equivalent intake of some artificial sweeteners also can lead to hypertension. However, Stevia, a sweetener that is isolated from a Paraguayan plant, was shown in relevant literature to decrease blood pressure in both rat specimens and humans. The general purpose of this research project was to study the effect of Stevia and glucose on the expression of two key components of the renin-angiotensin-aldosterone system (RAAS): prorenin receptor (PRR) and angiotensin receptor type 1 (AT1). Increased expression of renin and angiotensin can lead to vasoconstriction and systemic hypertension. Their effects are mediated by their binding to PRR and AT1. Therefore, decreases in the expression of these receptor proteins can result in lowered blood pressure. Rats were fed diets supplemented with glucose, saccharin, or Stevia over a six-week period and the kidneys were obtained. qPCR designs were developed to measure the relative amounts of PRR receptor and AT1 receptor. The methods had efficiencies greater than 97% and gave reproducible results. Then the developed methods were used to measure the expression of AT1 and PRR in the different rat kidney samples. A general increase in the relative expression of both AT1 and PRR in the glucose diet was observed. However, this increase and all other differences in the test groups were not significantly different than the control group. These results suggest no differences in AT1 or PRR expression for different sweetener diets.

Introduction




Figure 1. Simplified Diagram of Renin-Angiotensin System

- The renin-angiotensin-aldosterone system is located primarily in the kidneys
- It is important in regulating blood pressure and blood volume.
- Renin and angiotensin are the major proteins in this system; they bind to prorenin receptor (PRR) and angiotensin II type 1 receptor (AT1).
- Binding to these receptors lead to local vasoconstriction and hypertension.
- Hypertension is a serious health problem in the United States, more than 30% of adults have it.²
- AT1 expression is increased during hypertension.¹
- In rats, and increase in fructose & sucrose intake correlated with an increase in AT1 expression and hypertension.^{3,4,5}
- In both rats and humans, Stevia was shown to decrease blood pressure.^{6,7}
- Less information is known about the effects of sweeteners on prorenin receptor

Objectives

- To purify mRNA from the rat kidneys and assess its quality
- To design and test the efficiency of qPCR primers and probes to measure the expression of prorenin receptor (PRR), angiotensin II type 1 receptor (AT1), and GAPDH (endogenous control)
- To measure the expression of AT1 and PRR relative to GAPDH in the control and experimental rat kidneys

Experimental Design

Four groups, four male WKY rats per group
Fed the following liquid diets for 6 weeks:

Standard- liquid osmolite diet

Glucose- liquid osmolite & glucose

Saccharin- liquid osmolite & saccharin

Stevia- liquid osmolite & stevia


Mixed Diet - liquid osmolite (3 weeks), glucose (1 week), stevia (2 weeks) (2 kidneys)

Measured the relative expression of AT1 and PRR

Table 1. Primers and Probes Designed for AT1, PRR, and GAPDH

	Angiotensin II Receptor type 1 (AT1)	Prorenin Receptor (PRR)	Glyceraldehyde 3-phosphate dehydrogenase (GAPDH)
Forward	5' GCTGGCAITTTGTCTGGATAAAT 3'	5' TCGCTCCCCCTCAATTCCT 3'	5' ACGGGAAACCCATCACCAT 3'
Reverse	5' GGGTTGAGTTGGTCTCAGACACT 3'	5' CAGCACTGTCAGTTCAGAAAGAA 3'	5' CCAGCATCACCCCAITTTGA 3'
Probe	5' AGTGATCACCAGGTCAGT 3'	5' GGAATAATGAAGTTGACCTGC 3'	5' TTCCAGGAGCGAGATC 3'

Methodology



Forward Primer:
5' TCGCTCCC CTCATTCCT T 3'
(gagcagggg gagttaagag actcactctt attactt... 5')
19801 CTCCTCCCC CTCATTCCT TgagtAGGAA TAATGAA

Probe 5' GGAA TAATGAA GTTACCTG 3'
(catctctt attactt... 5')
CTGTGACCTG CcattTCCTT CTGAACCTGCA AGTCTGcat gata...
Reverse Primer 3' AAGAAA GACTTGACOT TCACGAC 5'

Forward Primer:
5' TCGCTCCC CTCATTCCT T 3' 5' GGAA TAATGAA GTTACCTG 3'
cDNA (3' gagcagggg gagttaagag actcactctt attacttcaa ctggacgata aagaaagact tgacttcaac gactact... 5')
5' TCGCTCCC CTCATTCCT TgagtAGGAA TAATGAA GTTACCTG CcattTCCTG ACTGCAAGT CTGcatga... 3'
Reverse Primer: 3' A AAGAAAGACT TGACCTCAC GAC 5'

Figure 2. Prorenin Receptor System Design. Crossing the exon junction ensures that the primers and probes are specific for only the mRNA of interest




Figure 3. Diagram of qPCR Theory. As polymerization occurs, the Fluorophore is cleaved from the probe and emits measurable fluorescence




Figure 4. Example Amplification Plot comparing the threshold cycles (C_T) of control and experimental targets. The C_T is the cycle where the fluorescence reaches above baseline fluorescence, generally during the exponential phase of the curve.

- $\Delta C_T = C_T (\text{target}) - C_T (\text{control} \rightarrow \text{GAPDH})$
- $\Delta \Delta C_T = \Delta C_T - \Delta C_T (\text{Ref} \rightarrow \text{E2})$
- $\text{RQ} = 2^{-(\Delta \Delta C_T)}$

Conclusions

- Efficient qPCR systems were established for AT1, PRR, and GAPDH.
- RNA was isolated with consistent purity despite a variance in concentration.
- No significant differences were found among sample groups, though a slight increase in the relative quantity of glucose was seen for both targets.
- The lack of significant differences between diet groups was probably a result of high variance among biological replicates and small sample sizes.

Results

RNA Purification

Table 2. RNA sample concentrations & purity

Sample	Conc (ng/uL)	A_{260}/A_{280}
E1	592.7	2.01
E2	111.9	1.97
E3	596.7	2.04
E4	439.7	2.02





Figure 5. 15% denaturing gel demonstrating the separation of short and long RNAs in the mirVana procedure

Efficiencies of RT-qPCR Methods

Efficiency Curve for AT1



Efficiency Curve for PRR




Figure 6. Standard curve to determine AT1 system efficiency. System was 97% efficient.

Figure 7. Standard curve to determine PRR system efficiency. System was 98% efficient.

Relative Expressions of AT1 and PRR mRNAs

Relative Expression of AT1 for Different Diets




Figure 7. Relative AT1 mRNA quantity for each diet group

Relative Expression of PRR for Different Diets




Figure 8. Relative PRR mRNA quantity for each diet group

References

- Contreras, F., Parte, A., Cabrera, J., Ospino, N., Hernandez, R., Hernandez, J., Lezama, E., & Velasco, M. (2002). Functional aspects of the renin-angiotensin-aldosterone system and angiotensin II receptor at1 blockers in arterial hypertension. *International Congress Series*, 1237(July), 87-98.
- Center for disease control and prevention: *Fastfacts hypertension*. (2013). Retrieved from <http://www.cdc.gov/nchs/fastats/hypertens.htm>
- Iyer, S. N., Raizadaz, M. K., & Katovich, M. J. (1996). AT1 receptor density changes during development of hypertension in hyperinsulinemic rats. *Clin and Exper. Hypertension*, 18(6), 793-810.
- Coelho, M. S., Lopes, K. L., Freitas, R. A., Barbosa, E. O., Bergamaschi, C. T., Campos, R. R., Casarini, D. E., & Carmona, A. K. (2010). High sucrose intake in rats is associated with increased ace2 and angiotensin-(1-7) levels in the adipose tissue. *Regulatory Peptides*, 162(1-3), 61-67.
- Shinozaki, K., Ayajiki, K., Nishio, Y., Sugaya, T., Kashiwagi, K., & Okamura, T. (2004). Evidence for a Causal Role of the Renin-Angiotensin System in Vascular Dysfunction Associated With Insulin Resistance Hypertension. 43:255-262
- Melis, M. S. (1995). Chronic administration of aqueous extract of stevia rebaudiana in rats: renal effects. *Journal of Ethnopharmacology*, 47(3), 129-134.
- Hsieh, M. H., Chan, P., Sue, Y., Liang, T. H., Huang, T., Tomlinson, B., Sum Chow, M. S., & Kao, P. (2003). Efficacy and tolerability of oral stevioside in patients with mild essential hypertension: A two-year, randomized, placebo-controlled study. *Clinical Therapeutics*, 25(11), 2797-2808.

Acknowledgements

Undergraduate Research Center Grant
Honors Program Grant
Dr. Theresa Salerno- Mentor
Dr. James Rife- Tissue Collection
Natalie Young – Research partner

Department of Chemistry & Geology
Dr. Penny Knoblich
Dr. David Bisonnette
Dr. Mary Hadley
& their research students