



Minnesota State University, Mankato
Cornerstone: A Collection of Scholarly
and Creative Works for Minnesota
State University, Mankato

All Graduate Theses, Dissertations, and Other
Capstone Projects

Graduate Theses, Dissertations, and Other
Capstone Projects

2014

Usage of the Alcohol Use Disorder Identification Test (AUDIT) With Hospital Inpatients

Jeni Kolstad
Minnesota State University, Mankato

Follow this and additional works at: <https://cornerstone.lib.mnsu.edu/etds>



Part of the [Clinical and Medical Social Work Commons](#), [Social Work Commons](#), and the [Substance Abuse and Addiction Commons](#)

Recommended Citation

Kolstad, J. (2014). Usage of the Alcohol Use Disorder Identification Test (AUDIT) with hospital inpatients [Master's capstone project, Minnesota State University, Mankato]. Cornerstone: A Collection of Scholarly and Creative Works for Minnesota State University, Mankato. <https://cornerstone.lib.mnsu.edu/etds/882/>

This Other Capstone Project is brought to you for free and open access by the Graduate Theses, Dissertations, and Other Capstone Projects at Cornerstone: A Collection of Scholarly and Creative Works for Minnesota State University, Mankato. It has been accepted for inclusion in All Graduate Theses, Dissertations, and Other Capstone Projects by an authorized administrator of Cornerstone: A Collection of Scholarly and Creative Works for Minnesota State University, Mankato.



MINNESOTA STATE UNIVERSITY MANKATO

Usage of the Alcohol Use Disorder Identification Test (AUDIT) With Hospital Inpatients

Jeni Kolstad, LSW

MSW Candidate, Minnesota State University, Mankato
Practicum Agency: Mayo Clinic Health System – Mankato Emergency Department
Field Instructor: Kelly Duloher, LICSW
Academic Advisor: David Beimers, PhD



Purpose

This project investigated whether the AUDIT was being used as the World Health Organization (WHO) recommends within Mayo Clinic Health System – Mankato (MCHS). The goals of the study were to describe the Chemical Dependency (CD) process and to determine the role of the AUDIT in the CD consult and assessment process. Recommendations were made to enhance the process of identifying, assessing, and intervening with patients with alcohol use problems.

Literature Review

Alcohol is the most frequently used psychoactive drug.¹ The AUDIT is one screening that can be used in a health care setting which can detect harmful and hazardous alcohol use and alcohol dependence before any physical or social problems occur.^{2,3,4}

The AUDIT is simple to use, standardized, and administered with little cost and time.^{2,4,5} It has been shown to have high internal consistency, reliability, and specificity.^{2,6} Most studies agree that a score of eight or more equals a positive screen, which would require further intervention.^{2,3,4,7,8} By using the AUDIT scoring system, brief interventions can be administered and can reduce alcohol consumption by up to 50% following a hospital inpatient stay.^{1,2} For patients scoring 20 or more a brief intervention is not appropriate and the patient should be referred to a program specializing in alcohol use disorders.⁷

References

- Shorie, S., Conigrave, K.M., Proude, E.M., & Haber, P.S. (2007). Detection of and intervention for excessive alcohol and tobacco use among adult hospital in-patients. *Drug and Alcohol Review*, 26, 127-133.
- Barbor, T.F., Higgins-Biddle, J.C., Saunders, J.B., & Monterio, M.G. (2001). The alcohol use disorders identification test: Guidelines for use in primary care. (Document No. WHO/MSD/MSB/01.6a).
- Chen, C.H., Chen, W.J., & Cheng, A.T.A. (2005). New approach to the validity of the alcohol use disorders identification test: Stratum-specific likelihood ratios analysis. *Alcoholism: Clinical and Experimental Research*, 29(4), 602-608.
- Conigrave, K.M., Saunders, J.B., & Reznik, R.B. (1995a). Predictive capacity of the AUDIT questionnaire for alcohol-related harm. *Addiction*, 90, 1479-1485.
- Berner, M.M., Krston, L., Bentele, M., & Harter, M. (2007). The alcohol use disorders identification test for detecting at-risk drinking: A systematic review and meta-analysis. *Journal of Studies on Alcohol and Drugs*, 68, 461-473.
- Conigrave, K.M., Hall, W.D., & Saunders, J.B. (1995b). The AUDIT questionnaire: Choosing a cut-off score. *Addiction*, 90, 1349-1356.
- Saitz, R. (2013). Screening for unhealthy use of alcohol and other drugs. *UpToDate*.
- Selin, K.H. (2006). Alcohol use disorder identification test (AUDIT): What does it screen? Performance of the AUDIT against four different criteria in a Swedish population sample. *Substance Use and Misuse*, 41, 1881-1899.

Methodology

This project had two distinct parts. Part I involved oral, semi-structured interviews with MCHS staff about the CD consult and assessment process. Ordering doctors and consulted social workers were asked four open-ended questions about their perceptions of the CD process, if they use the AUDIT in determining a need for consult or use the AUDIT in assessing patients, and any recommended changes they have for the process.

Part II asked the social workers to document on a score card the AUDIT score of the patient and the intervention/recommendation provided to the patient.

Results - Interviews

Staff Interviews – 11 Interviews Completed (7 Social Workers & 4 Doctors)


- Only 1 staff used the AUDIT when assessing patients for chemical use.
- All social workers interviewed (7) wanted more consistency with the ordering, assessing, and staffing for completion of CD consults.
- 71% of social workers wanted the ability to screen out inappropriate CD consults.
- All doctors interviewed (4) used the social worker's recommended interventions for their patients.
- Doctors wanted to see consults done in a timelier manner, better communication with the doctors, and more education on CD use provided to patients.

Results – AUDIT Scorecard

Audit Score and Drinking Risk Level	# of Assessments Completed	Recommended intervention by World Health Organization	% Followed Recommended Intervention
1-7: Low risk drinking	15	Education	40%
8-15: Use in excess of low-risk	7	Simple advice	0%
16-19: Harmful and hazardous drinking	4	Simple advice, plus brief counseling and continued monitoring	0%
20-40: Alcohol dependence	8	Referral to specialist for diagnostic evaluation and treatment	88%
34 total AUDIT scores were collected with recommendations following AUDIT guidelines 38% of the time.			

Recommendations

The AUDIT tool can alleviate many of the concerns that both social workers and doctors have about the consult process. The AUDIT provides social workers with the ability to be more consistent in assessing patients and to screen out CD consults. Because the AUDIT can be completed in less than 10 minutes, it allows social workers to be more timely in seeing patients.



Flow Chart

As part of the assessment, the social worker should administer the AUDIT on every consult. The AUDIT score combined with clinical judgment should be used to determine if a full CD assessment is needed. The purpose of the full assessment is to determine if the patient needs services beyond what can be offered within the hospital. Brief interventions should be done with patients not needing additional services.

Brief Interventions

Elements of brief interventions include:

- Provide feedback about the results of the screening.
- Educate about low-risk levels of drinking and hazards of exceeding them.
- Encourage patients to adhere to the guidelines.
- Simple advice may be needed for people who score between an 8 and 15 on the AUDIT. This includes discussing the risks of drinking in excess and the need to cut down or stop drinking all together.
- Work with patients on creating a goal. Patients reluctant to create a goal may need more than a brief intervention.